

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I being given ZYPREXA?

ZYPREXA contains the active ingredient olanzapine. ZYPREXA is used to help correct chemical imbalances in the brain, which may cause mental illness.

For more information, see Section [1. Why am I being given ZYPREXA?](#) in the full CMI.

2. What should I know before being given ZYPREXA?

Do not use if you have ever had an allergic reaction to olanzapine or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before being given ZYPREXA?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with ZYPREXA and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How am I being given ZYPREXA?

Your doctor will decide how much ZYPREXA IM you need and how long you need it for. The dosage you receive will usually be 10mg in one injection. Up to three injections in 24 hours may be given.

A lower starting dose may be prescribed for elderly patients over the age of 65 years.

More instructions can be found in Section [4. How am I being ZYPREXA?](#) in the full CMI.

5. What should I know while having ZYPREXA?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using ZYPREXA• Tell your doctor if you become pregnant while taking ZYPREXA• Keep all your doctor's appointments so that your progress can be checked
Things you should not do	<ul style="list-style-type: none">• Do not stop taking your medicine, or change the dosage, without first checking with your doctor.• Do not give this medicine to anyone else, even if they have the same condition as you
Driving or using machines	<ul style="list-style-type: none">• Be careful before you drive or use any machines or tools until you know how ZYPREXA affects you. ZYPREXA may cause drowsiness in some people
Drinking alcohol	<ul style="list-style-type: none">• Tell your doctor if you drink alcohol. The effects of alcohol could be made worse while taking ZYPREXA
Looking after your medicine	<ul style="list-style-type: none">• ZYPREXA IM injection vials should be stored below 25° C• The vials are usually stored in the hospital ward or in the pharmacy• Keep this medicine where children cannot reach it

For more information, see Section [5. What should I know while having ZYPREXA?](#) in the full CMI.

6. Are there any side effects?

Like other medicines, ZYPREXA may cause some unwanted side effects. These are likely to vary from patient to patient, although not everybody gets them. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need urgent medical attention.

Tell your doctor if you notice anything unusual or if you are concerned about any aspect of your health, even if you think the problems are not connected with this medicine and are not referred to in this leaflet.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

ZYPREXA®IM

Active ingredient(s): Olanzapine

Consumer Medicine Information (CMI)

This leaflet provides important information about using ZYPREXA. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using ZYPREXA.**

Where to find information in this leaflet:

- [1. Why am I being given ZYPREXA?](#)
- [2. What should I know before being given ZYPREXA?](#)
- [3. What if I am taking other medicines?](#)
- [4. How am I being given ZYPREXA?](#)
- [5. What should I know while having ZYPREXA?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am being given ZYPREXA?

ZYPREXA contains the active ingredient olanzapine.

Olanzapine belongs to a group of medicines called antipsychotics. It helps to correct chemical imbalances in the brain, which may cause mental illness.

ZYPREXA IM injection is used for the rapid control of agitation and disturbed behaviors in patients with schizophrenia and related psychoses and in patients with acute mania associated with Bipolar I Disorder.

Schizophrenia is a mental illness with disturbances in thinking, feelings and behavior. Bipolar I Disorder is a mental illness with symptoms such as feeling "high", having excessive amounts of energy, needing much less sleep than usual, talking very quickly with racing ideas and sometimes severe irritability.

ZYPREXA IM injection is given when treatment with ZYPREXA tablets is not appropriate. Your doctor will change your treatment to ZYPREXA tablets or ZYPREXA Zydys wafers as soon as practicable.

Your doctor may have prescribed ZYPREXA IM for another reason.

Ask your doctor if you have any questions about why ZYPREXA has been prescribed for you.

This medicine is available only with a doctor's prescription.

ZYPREXA is not recommended for use in children under the age of 18 years as there is not enough information on its effects in this age group.

2. What should I know before being given ZYPREXA?

Warnings

Do not use ZYPREXA if:

- you are allergic to olanzapine, or any of the ingredients listed at the end of this leaflet.
- Always check the ingredients to make sure you can use this medicine.
- Signs of an allergic reaction may include a skin rash, itching, shortness of breath or swelling of the face, lips or tongue.
- the packaging is torn or shows signs of tampering, or the injection vials do not look quite right.
- the expiry date on the pack has passed. If you take this medicine after the expiry date has passed it may not work as well.

Use of ZYPREXA IM is not recommended simultaneously with a benzodiazepine medication by injection (to treat a range of conditions such as anxiety, nervousness, panic attacks, seizures, muscle spasm and insomnia).

If you are not sure whether you should have ZYPREXA, talk to your doctor or pharmacist.

Check with your doctor if you:

- **have had an allergic reaction to any medicine which you have taken previously to treat your current condition.**
- **have any other medical conditions, especially the following:**
 - tumor of the pituitary gland (a small gland at the base of the brain)
 - disease of the blood or bone marrow with a reduced number of white or red blood cells
 - disease of the blood vessels of the brain, including stroke
 - prostate problems
 - kidney or liver disease
 - high blood sugar, diabetes or a family history of diabetes
 - breast cancer or a family history of breast cancer
 - paralytic ileus, a condition where the small bowel does not work properly
 - epilepsy, seizures or fits
 - glaucoma, a condition in which there is usually a buildup of fluid in the eye
 - heart disease, including irregular heart rhythm
 - neuroleptic malignant syndrome, a reaction to some medicines with a sudden increase in body temperature, extremely high blood pressure and severe convulsions

- tardive dyskinesia, a reaction to some medicines with uncontrollable twitching or jerking movements of the arms and legs.
- sleep apnea, a sleep disorder where a person has pauses in breathing or periods of shallow breathing during sleep.

Tell your doctor if

- you will be in a hot environment or do a lot of vigorous exercise. ZYPREXA may make you sweat less, causing your body to overheat.
- you smoke. Smoking may affect ZYPREXA or may affect how it works.
- you are elderly with dementia-related psychosis
- you have previously had a stroke or transient ischaemic attack (mini-stroke) or if you have high blood pressure.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Like most antipsychotic medicines, ZYPREXA is not recommended for use during pregnancy.

Newborn babies of mothers taking antipsychotic drugs (including ZYPREXA) during the last trimester of pregnancy are at risk of experiencing extrapyramidal neurological disturbances and/or withdrawal symptoms following delivery. These may include, but are not limited to agitation, tremor, muscle stiffness or weakness, drowsiness, feeding problems, and breathing difficulty. If there is a need to consider ZYPREXA during your pregnancy, your doctor will discuss with you the benefits and risks of using it.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

It is recommended that you do not breast-feed while taking ZYPREXA.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may be affected by ZYPREXA or may affect how it works. These include:

- medicines used to treat a fast or irregular heartbeat (arrhythmia).
- medicines taken for anxiety or to help you sleep.
- medicines taken for depression
- carbamazepine, a medicine used for mood stabilization and to treat epilepsy

- other centrally acting medicines (eg. tranquilizers or strong painkillers)
- ciprofloxacin, a medicine used to treat bacterial infections
- medicines that lower blood pressure
- medicines used for Parkinson's disease
- medicines that can change the heart's electrical activity or make it more likely to change.

Your doctor or pharmacist has more information on medicines to be careful with or avoid while taking ZYPREXA.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect ZYPREXA.

4. How am I being given ZYPREXA?

How much to take

Your doctor will decide how much ZYPREXA you need and how long you need it for.

The dosage you receive will usually be 10mg in one injection. Up to three injections in 24 hours may be given.

A lower starting dose may be prescribed for elderly patients over the age of 65 years.

How it is given

ZYPREXA is injected into your muscle. Your doctor or nurse will inject ZYPREXA for you.

If you are given too much ZYPREXA

As ZYPREXA IM is given to you under the supervision of your doctor, it is unlikely that you will receive too much. However, if you experience any side effects after being given ZYPREXA, tell your doctor immediately or go to Accident and Emergency at your nearest hospital.

The most common signs of too much ZYPREXA are fast heartbeat, agitation/aggression, difficulty speaking, uncontrollable movements and sleepiness (sedation).

You should immediately:

- phone the Poisons Information Centre (by calling 13 11 26), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while having ZYPREXA?

Things you should do

Tell all doctors, dentists and pharmacists who are treating you that you are having ZYPREXA.

If you are about to be started on any new medicine, remind your doctor, dentist or pharmacist that you are having ZYPREXA.

Tell your doctor immediately if you become pregnant while taking ZYPREXA.

Keep all of your doctor's appointments so that your progress can be checked.

- Your doctor should monitor your weight while you are taking ZYPREXA.
- Patients with diabetes or who have a higher chance of developing diabetes should have their blood sugar checked often.
- Your doctor may request you have a blood test from time to time to monitor your cholesterol levels.
- If you are over 65, your doctor may measure your blood pressure from time to time.

Tell your doctor if you are female and your monthly periods are absent for six months or more.

All thoughts of suicide or violence must be taken seriously. Talk to your doctor or mental health professional if you have thoughts about death or suicide; or thoughts or talk about self-harm, or harm to others.

These may be signs of changes or worsening in your mental illness.

Things you should not do

Do not stop taking ZYPREXA or lower the dosage, even if you are feeling better, without checking with your doctor.

Do not give ZYPREXA to anyone else, even if their symptoms seem similar or they have the same condition as you. Your doctor has prescribed ZYPREXA for you and your condition.

Things you should be careful of

If ZYPREXA makes you feel light-headed, dizzy or faint, be careful when getting up from a sitting or lying position. Getting up slowly may help.

If outdoors, wear protective clothing and use at least a 30+ sunscreen.

ZYPREXA may cause your skin to be much more sensitive to sunlight than it is normally. Exposure to sunlight may cause a skin rash, itching, redness, or severe sunburn.

If your skin does appear to be burning, tell your doctor.

Make sure you keep cool in hot weather and keep warm in cool weather. ZYPREXA may affect the way your body reacts to temperature changes.

Antipsychotics have the potential to cause cardiac complications and sudden cardiac death.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how ZYPREXA affects you.

ZYPREXA may cause dizziness in some people.

Drinking alcohol

Tell your doctor if you drink alcohol.

The effects of alcohol could be made worse while taking ZYPREXA.

Your doctor may suggest you avoid alcohol while you are being treated with ZYPREXA.

Looking after your medicine

ZYPREXA IM Injection vials should be stored below 25° C.

The vials are usually stored in the hospital ward or in the pharmacy.

Keep it where young children cannot reach it.

There will be an expiry date (month, year) on your ZYPREXA container.

The medicine should not be taken after this date because it may have lost some of its strength.

Getting rid of any unwanted medicine

If your doctor tells you to stop taking ZYPREXA or you find that the tablets or wafers have passed their expiry date, please return any left over tablets or wafers to your pharmacist.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Like other medicines, ZYPREXA may cause some unwanted side effects. These are likely to vary from patient to patient. Some side effects may be related to the dose of ZYPREXA. Accordingly, it is important that you tell your doctor as soon as possible about any unwanted effects. Your doctor may then decide to adjust the dose of ZYPREXA you are taking.

Some people may feel dizzy in the early stages of treatment, especially when getting up from a lying or sitting position. This side effect usually passes after taking ZYPREXA for a few days.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none"> • drowsiness • unusual tiredness or weakness • fever • restlessness or difficulty sitting still • increased appetite, weight gain • constipation, bloating • dry mouth • swelling of your hands, feet and ankles • aching joints • nose bleeds • changes to heart rate, such as fast heartbeat or slow heartbeat • low blood pressure • dizziness, confusion, forgetfulness • speech disorder • sleepwalking • sleep eating 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none"> • symptoms of sunburn (such as redness, itching, swelling or blistering of the skin) which occur more quickly than normal • rash or allergic reaction • slow heartbeat • changes in sexual functioning or sex drive in men or women • prolonged and/or painful erection • unusual secretion of breast milk • breast enlargement in men or women • symptoms of high sugar levels in the blood (including passing large amounts of urine, excessive thirst, having a dry mouth and skin and weakness) These may indicate the onset or worsening of diabetes • reaction following abrupt discontinuation (profuse sweating, nausea or vomiting) • absence of menstrual periods and changes in the regularity of menstrual periods • involuntary passing of urine or difficulty in initiating urination • unusual hair loss or thinning 	<p>Tell your doctor as soon as possible if you notice any of these serious side effects. You may need medical attention.</p>

Very serious side effects

Very serious side effects	What to do
<ul style="list-style-type: none"> • Sudden signs of an allergic reaction such as a skin rash, itching, shortness of breath or swelling of the face, lips or tongue. • frequent infections such as fever, severe chills, sore throat or mouth ulcers • bleeding or bruising more easily than normal. • painful swollen leg, chest pain, or shortness of breath as these can be signs of blood clots in the lungs or legs • seizures, fits or convulsions • yellowing of the skin and/or eyes • nausea, vomiting, loss of appetite, generally feeling unwell, fever, itching • severe upper stomach pain often with nausea and vomiting (inflammation of the pancreas) • worm-like movements of the tongue, or other uncontrolled movements of the tongue, mouth, cheeks, or jaw which may progress to the arms and legs • sudden increase in body temperature, sweating, fast heartbeat, muscle stiffness, high blood pressure and convulsions • sharp chest pain, coughing of blood, or sudden shortness of breath • pain/tenderness in the calf muscle area • muscle pain, muscle weakness and brown urine • heart attack • heart palpitations and dizziness, which may lead to collapse. • fast breathing, shortness of breath, fever with chills, feeling tired or weak, chest pain while coughing, fast heartbeat. You may have pneumonia. • fever or swollen glands, especially if they occur together with or shortly after a skin rash 	<p>Call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects. You may need urgent medical attention or hospitalization.</p>

The following additional side effects may occur in some groups of people taking ZYPREXA:

Elderly patients with dementia-related psychosis

- unusual manner of walking
- falls
- pneumonia
- involuntary passing of urine
- Stroke
- transient ischemic attack – symptoms maybe, but not limited to, paralysis in face, arm or leg.

Parkinson's disease psychosis

Some patients with Parkinson's disease may hallucinate (see, feel or hear things that are not there) or develop worsening symptoms of Parkinson's disease.

In combination with lithium or valproate

Patients with bipolar mania taking ZYPREXA in combination with lithium or valproate may notice the following additional side effects:

- tremors
- speech disorder

Tell your doctor if you notice anything unusual or if you are concerned about any aspect of your health, even if you think the problems are not connected with this medicine and are not referred to in this leaflet.

Some side effects, such as changes to liver function, blood cell counts, cholesterol or triglycerides can occur. These can only be found when your doctor does tests from time to time to check your progress.

Do not be alarmed by this list of side effects. Most of these side effects are very rare. You may not experience any of them. Your doctor has weighed the risks of using this medicine against the benefits they expect it will have for you.

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What ZYPREXA IM contains

Active ingredient (main ingredient)	Olanzapine
Other ingredients (inactive ingredients)	Lactose monohydrate Tartaric acid
Potential allergens	None

ZYPREXA IM injection comes as a yellow powder in a clear glass vial. A vial of ZYPREXA IM injection can provide you with 10 mg of olanzapine. (AUST R 76867)

Who distributes ZYPREXA IM

Pharmaco (Australia) Ltd
Level 9, Tower A
Zenith Towers
821 Pacific Highway
Chatswood, NSW 2067
Australia
Phone: 1800 201 564

Under license of CHEPLAPHARM Schweiz GmbH,
Switzerland

This leaflet was prepared in February 2025.